


## Appendix C

Plant Molecular Biology  
39 (5): 891-906, March 1999  
Copyright © 1999 Kluwer Academic Publishers  
All rights reserved



RECEIVED

AUG 01 2002

TECH CENTER 1600/2900

## **$\Delta$ 7-Sterol-C5-desaturase: molecular characterization and functional expression of wild-type and mutant alleles**

**Tania Husselstein**

*Institut de Biologie Moléculaire des Plantes, Département d'Enzymologie Cellulaire et Moléculaire, UPR 406 du CNRS, 28 rue Goethe, 67083 Strasbourg Cédex, France*

**Hubert Schaller**

*Institut de Biologie Moléculaire des Plantes, Département d'Enzymologie Cellulaire et Moléculaire, UPR 406 du CNRS, 28 rue Goethe, 67083 Strasbourg Cédex, France*

**Daniel Gachotte**

*Present address: Department of Biology, Indiana University-Purdue University at Indianapolis, 723 W. Michigan Street, Indianapolis, IN 46202, USA*

**Pierre Benveniste**

*Institut de Biologie Moléculaire des Plantes, Département d'Enzymologie Cellulaire et Moléculaire, UPR 406 du CNRS, 28 rue Goethe, 67083 Strasbourg Cédex, France*

### **Abstract**

An *Arabidopsis thaliana* recessive monogenic mutant (*ste1-1*) presenting a deficiency of the  $\Delta$ 7-sterol-C5(6)-desaturase step in the sterol pathway has been reported previously [12]. To further characterize *ste1-1*, *Arabidopsis*, *Nicotiana tabacum* and *Homo sapiens* cDNAs encoding  $\Delta$ 7-sterol-C5(6)-desaturases were isolated and identified on the basis of their ability to restore ergosterol synthesis in *erg3*, a yeast null mutant whose gene encoding the  $\Delta$ 7-sterol-C5(6)-desaturase was disrupted. Overexpression of the *Arabidopsis* cDNA driven by a 35S promoter in transgenic *ste1-1* plants led to full complementation of the mutant. This result demonstrates that STE1 was the impaired component in the desaturation system. Four independent reverse transcriptions of *ste1-1* RNA followed by polymerase chain reactions (RT-PCRs), yielded a single product. Alignment of the wild-type ORF with the RT-PCR derived *ste1-1* ORF revealed a single amino acid substitution: Thr-114 in the wild-type is changed to Ile in *ste1-1*. Expression in *erg3* resulted in a 6-fold lowered efficiency of the *ste1-1* ORF in complementing the yeast biosynthetic pathway when compared to the wild-type ORF. The presence of this mutation in the mutant *ste1-1* genomic sequence (and no additional modification between *ste1-1* and wild-type genes) demonstrates that the change of the Thr-114 to Ile is necessary and sufficient to create the leaky allele *ste1-1*. The occurrence of a hydroxylated amino acid (Thr or Ser) at the position corresponding to Thr-114 in the five  $\Delta$ 7-sterol-C5(6)-desaturases identified so far suggests that this amino acid is important for normal enzymatic function.

### **Keywords**

sterol,  $\Delta$ 7-sterol-C5(6)-desaturase, deficient mutant, complementation, *Arabidopsis*

COPY OF PAPERS  
ORIGINALLY FILED

**Article ID: 200124**

**SearchPLUS**

**Open SearchPLUS Navigator**

Use the pop-up window to navigate search results and find highlighted terms in this document.

☐ Hide SearchPLUS

☐ Disable Highlights

**Reload this page**